

A drug-free option for pain relief during labor



Start the TENS unit in early labor for best results

GIVE US A RING WITH ANY **OUESTIONS** 844-997-1431


1. Wipe the areas of your back where the pads will go with an alcohol wipe

It's safe to practice using a tens after 37 weeks pregnancy

- 2. Partners: place electrode pads as shown on the next page. 1 inch on either side of the spine. Upper pads to go underneath the where bra strap would be with wires facing down. Lower pads to line up with Gluteal cleft or "butt crack" wires facing down. See image
- 3. Insert lead wire pins into stimulating pads (see image)
- 4. Make sure the tens unit is turned OFF
- 5. Insert the jack into the socket of the OBI TENS unit
- 6. Give the TENS unit to the laboring mother.

YOU'VE GOT

- 7. Turn the unit on and increase the plus button up until you feel a mild tickling sensation. Increase intensity to a comfortable level.
- 9. Press the BOOST during a contraction! Use the BOOST mode DURING a contraction and switch back to burst mode BETWEEN contractions



- 11. Replace reusable electrodes pads on the special sheet they came on and store in the ziploc bag for future use
- 12. If used and stored correctly, the pads will stick for 10-15 times







How to set up and operate the OBI TENS unit



Attach two (upper) electrodes 1 inch from spine and underneath where "bra strap" would be.

Attach two (lower) electrodes 1 inch from spine and bottom of pads to line up with gluteal cleft or better known as "butt crack"

Attach the red connector wires to one set of electrodes and the white connector wire to the other set of electrodes. (top or botton sets of electrodes, it doesnt matter.



Make sure tens unit is in **off** position. Plug lead wire into BOTTOM of OBI tens unit



Turn on TENS unit. Control the intensity of the pads on the OBI tens unit by the up and down arrows on the tens unit. Use the boost button when you have a contraction for added pain relief

Contraindications

- TENS must NOT be used before the 37th week of pregnancy.
- If you are pregnant, do NOT place the electrodes over your abdomen.
- Do NOT use if you have a pacemaker or any other implanted electrical device.
- Check with your healthcare provider before using if you suffer from a heart condition, epilepsy, undiagnosed pain, or have any metal implants.
- Do NOT use if you experience an allergic reaction to the
- electrodes.